

# BEEF CO-BALANCER R-800 MINERAL

**TYPE B MEDICATED FEED**  
FOR MANUFACTURE OF TYPE C MEDICATED FEED FOR CATTLE

For improved feed efficiency in cattle being fed for slaughter in confinement. For increased rate of weight gain in cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT	
Monensin Sodium .....	800 g/ton
GUARANTEED ANALYSIS	
Crude Protein .....	15.5%
(This includes not more than 10% equivalent crude protein from non-protein nitrogen)	
Calcium, min. ....	26.5%
Calcium, max. ....	29.5%
Phosphorus, min. ....	0.3%
Salt, min. ....	7.6%
Salt, max. ....	9.1%
Magnesium, min. ....	0.3%
Potassium, min. ....	1.4%
Selenium, min. ....	9.0 ppm
Vitamin A, min. ....	90,000 IU/lb
Vitamin D3, min. ....	20,000 IU/lb
Vitamin E, min. ....	200 IU/lb

**INGREDIENTS**

Calcium Carbonate, Plant Protein Products, Salt, Urea, Potassium Chloride, Monocalcium and Dicalcium Phosphates, Zinc Sulfate, Magnesium Oxide, Manganese Sulfate, Copper Sulfate, Sodium Selenite, dl-Alpha-Tocopheryl Acetate, Thiamine Mononitrate, Vitamin A Supplement, Vitamin D3 Supplement, Cobalt Sulfate, Ethylenediamine Dihydriodide, and Mineral Oil.

Take Time  Observe Directions on Back of Label

**CAUTION:** Not to exceed 0.3 ppm of selenium in total ration. The addition of higher levels of premix containing selenium is not permitted.

Manufactured For  
MIDWESTERN BIOAG  
BLUE MOUNDS, WI 53517  
NET WT 50 LB (22.67 kg)

**MIXING/FEEDING DIRECTIONS**

**\*\*DO NOT FEED UNDILUTED\*\***

For improved feed efficiency in cattle being fed in confinement for slaughter: Feed continuously in complete feed (5–40 g monensin /ton) at a rate of 50 to 480 mg of monensin per head per day as follows\*:

Desired Monensin Level g/ton (mg/lb)	Amount of Type B Feed to Add (lbs.)	Amount of non- medicated feed (lbs)	Feeding amount of finished feed (lb/hd/d)
5 (2.5)	12.5	1987.5	20.0 – 80.0
10 (5.0)	25.0	1975.0	10.0 – 40.0
20 (10.0)	50.0	1950.0	5.0 – 20.0
25 (12.5)	62.5	1937.5	4.0 – 16.0

\*The selenium concentration of this Type B Feed limits its inclusion rate to 62.5 lbs per ton of Type C Feed (25 g monensin/ton).

For increased rate of weight gain in cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers): Mix Type B medicated feed with non-medicated feed to a final concentration of 25 – 400 g monensin/ton and feed to provide 50 to 200 mg monensin per head per day. Mix and feed according to the table below. During the first 5 days of feeding, cattle should receive no more than 100 mg monensin per day contained in not less than 1 lb. of feed. Thereafter, feed to supply 50 to 200 mg monensin per head per day in not less than 1 lb. of feed or, after the fifth day feed at a rate of 400 mg per head every other day in not less than 2 lb. of feed.

Desired Monensin Level g/ton (mg/lb)	Amount of Type B Feed to Add (lbs.)	Amount of non- medicated feed (lbs)	Feeding amount of finished feed (lb/hd/d)
25 (12.5)	62.5	1937.5	4.0 – 16.0
100 (50)	125.0	1875.0	1.0 – 4.0
200 (100)	250.0	1750.0	1.0 – 2.0
300 (150)	375.0	1625.0	1.0 – 1.3
400 (200)	500.0	1500.0	1.0

**CAUTION:** Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by equines has been fatal. Monensin-medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Must be thoroughly mixed in feeds before use. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

**WARNING:** A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

**NOTICE:** Type C cattle feed containing 30 g monensin sodium or less per ton shall bear an expiration date of 30 days after the date of manufacture.

**Feed According to Your Consultant's Recommendations.**